



Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

62,31

Indexed



DESCRIPTIVE CIRCULAR

—OF—

FRUIT TREES, PLANTS,
VINES, ETC.

OFFERED AT



FOR THE

FALL OF 1893, AND SPRING OF 1894.

BY

H. F. HILLENMEYER.

111

We Employ No Agents, but Sell Direct to the Planter.



PRICE LIST

FOR THE SPRING OF 1894.

STANDARD APPLES.

	1	10	100	1000
Extra trees, 3 to 4 years, 5 to 7 feet.....	25	\$1 75	\$10 00	\$90 00
Good trees, 2 to 3 years, 5 to 6 feet.....	20	1 25	6 00	50 00

PEARS.

Standard or dwarf trees, very fine	35	3 00	25 00	
Light trees, 4½ to 5 feet.....	20	2 00	15 00	

PEACHES.

One year, extra, 5 to 7 feet.....	20	1 50	10 00	
One year, medium, 4 to 5 feet.....		1 25	6 00	
One year, light, 3 to 4 feet.....			3 00	

CHERRY.

Two year trees.....	30	2 50	16 00	
One year, or light two year.....	25	1 75	10 00	

PLUM.

Trees on plum roots, 4½ to 6 feet, very fine....	30	2 50	20 00	
--	----	------	-------	--

CURRENTS and GOOSEBERRIES.

Two year plants		50	4 00	
-----------------------	--	----	------	--

RASPBERRIES.

Black			1 00	7 50
Red.....			1 00	6 00

GRAPES—Large Stock.

Catawba and Cottage, two year.....	20	1 50	6 00	
Concord, Ives and Elvira, extra two year	15	1 00	4 00	25 00
Concord, Ives and Elvira, good 2 year	10	75	2 00	15 00
Reisling, Perkins' Brighton.....	20	1 25	8 00	
Martha, Noah, Pocklington, 2 year.....	20	1 50	8 00	
Norton, Delaware. Worden, 2 year.....	25	2 00		
Niagara, Woodruff, Wyoming, Empire State...	25	2 00		

PERSIMMON.

Two year, seedless.....	35	3 00		
From Seed.....	25	1 50		

STRAWBERRIES.

Special catalogue.....			50	3 50
------------------------	--	--	----	------

QUINCES.

Two year	25	2 00		
----------------	----	------	--	--

ASPARAGUS.**1 10 100 1000**

Colossal, 1 year.....			50	3 00
Colossal, 2 year, fine roots.....			75	5 00
Colossal, 3 year, extra.....			1 00	7 50

RHUBARB.

Victoria Linnaeus, divided crowns.....		50	3 00	
--	--	----	------	--

SAGE.

Fine Plants.....	5	40	3 00	
------------------	---	----	------	--

ORNAMENTAL TREES.

Ash, European, 7 to 10 feet.....	30	2 50		
Alder, English, 9 to 10 feet, fine.....	30	2 50		
Birch, European, white, 8 to 10 feet.....	30	2 50		
Catalpa, in variety, 9 to 10 feet.....	25	2 00	18 00	
Elms, in variety, 8 to 10 feet, stems 1½ to 1½ in.	30	2 50	20 00	
Liquidamber, 6 to 8 feet, fine.....	30	2 50		
Larch, 4 to 5 feet, fine.....	30	2 50		
Linden, European or American, 7 to 10 feet....	30	2 50		
Maple, Silver, 1½ inch stems.....	30	2 50	20 00	
Maple, Silver, 1¼ inch stems.....	25	2 00	15 00	
Maple, Sugar, 8 to 10 feet, fine.....	30	2 50	20 00	
Maple, Norway and Sycamore, 6 to 10 feet....	30	2 50		
Mountain Ash, 8 to 10 feet.....	30	2 50	18 00	
Magnolia, 5 to 7 feet.....	50	4 00		
Maiden Hair, Salisburia, 6 to 8 feet.....	30	2 50		
Mulberry, Russian.....	25	2 00		
Oak, English, 8 feet.....	30	2 50		
Poplar, Lombardy and Carolina, 10 to 14 feet..	30	2 50	20 00	
Poplar, Tulip, 8 to 10 feet.....	30	2 50	20 00	
Willow, Weeping, 6 to 8 feet.....	25	2 00		
Walnut English, 4 feet.....	25	2 00		
Hydrangea, Paniculata.....	25	2 00		

EVERGREENS.

Arbor Vitæ assorted, 3 to 4 feet.....	25	2 00	15 00	
Hemlock, 3 to 4 feet.....	30	2 50	20 00	
Juniper, Irish, 2 feet.....	35	3 00		
Retinospora, in variety.....	30	2 50		
Norway Spruce, 2 to 3 feet.....	30	2 50	18 00	
Norway Spruce, 3 to 4 feet.....	35	3 00	20 00	
Silver Fir, 3 to 4 feet.....	30	2 50	20 00	
White Pine, 5 feet, fine.....	35	3 00		
White Pine, 3 to 4 feet, fine.....	30	2 50		
Scotch and Austrian Pine.....	30	2 50		

SHRUBS--Hardy, Fine Plants in variety.

Althea, Deutzia, Calycanthus, Dogwood, Red- bud, Syringa, Wigelia, Tamerax, Pyrus- Japonica, Spirea, Lilac, Purple Fringe, Honeysuckle, Virginia Creeper, etc.....	20	1 50		
---	----	------	--	--

TO OUR PATRONS...

IN PRESENTING the seventeenth issue of this catalogue of nursery products, a feeling of thankfulness is entertained for the generous support extended by widely scattered patrons from the beginning of this enterprise. Identified for two generations with the fruit and nursery interests of Central Kentucky, it has been our effort to prosper by the success rather than the failure of our patrons, and as each year has seen an increase in our stock and a more widely extended trade, the hope is entertained that the effort has been successful.

The early summer of 1893 was very favorable for all kinds of stock, and the season later becoming very dry, every twig is ripened to the very tip, insuring trees likely to do well next year. A personal examination of our large and varied assortment is invited. Our location is so favorable that almost every thing used is grown here, and the propagation and distribution is all under our own personal supervision.

NO AGENTS.

We employ no agents, and never have. Our methods are incompatible. Trees will die, years of failure will come, and blight, frost and insect visitations will marr the brightest prospect. These cannot be obviated, even with the widest knowledge and most favorable locations, and we frankly admit that such is the case.

We make no claims for our goods except that, with diligent care, such as will achieve success in other callings, the outcome will be satisfactory. Agents promise so much more than is likely to be realized that we cannot assume the responsibility.

Many of our friends each year solicit orders and advance our interests simply through good will because of previous satisfactory dealings. These evidences of kindly interest are highly appreciated, and no opportunity of reciprocating will be lost.

A CHAT WITH OUR PATRONS.

To be successful with fruit and ornamental trees requires patient and intelligent labor and at least fairly rich, dry soil. The preparation should be thorough, the holes ample and the cultivation at least as good as that of corn. Small grain is ruinous to a young orchard, but any tilled crops may be advantageously planted till the ground is occupied by the trees. When trees are received they should be unpacked and planted at once, but if the soil is too wet or not prepared, they should be set in a trench until wanted. If received frozen, or the weather is inclement, trees should be left to thaw in the cellar. It will not injure them, as the roots of trees and plants in the open ground are frozen many times during the winter. A tree should be planted about as deep as it grew in the nursery, the roots spread in the natural way, and the branches either shortened one-half or the weaker ones entirely cut out. In digging, more or less roots are lost, and the head must be reduced in proportion. Small holes, a failure to prune, and the lack of cultivation, cause nine-tenths of the loss and dissatisfaction of tree planting. When trees die, the nurseryman generally gets the blame, but these causes are beyond his control. If you have not time to plant well, to prune and to cultivate, do not plant at all. You will lose money and temper, and the price of the goods will not compensate for your ill-will. A nurseryman is not benefitted by the patrons that fail. It is those that succeed, that have rich harvests, and great profits, that support him. Profit made by the loss of another cannot be long sustained.

Pear, Cherry and Plum trees should be headed low, and a wrapping of straw or paper around the stems will protect from heat, cold and rabbits. All trees planted near walls, buildings, etc., where the heat is intensified by radiation, should be thus protected. All the orchards planted here have been thus treated, with gratifying results. After two years' growth, the stems may annually in May be whitewashed to remove all scales, lichens and insects, answering a better purpose than more expensive applications of soda or potash. It is almost a waste of time and effort to replant vacancies in an old orchard.

The distance at which trees should be planted is dependent on the fertility of the soil. Peach and Pear should be from 16 to 20 feet in square; Plum and Cherry from 14 to 16 feet, and Apples from 28 to 32 feet; Currants and Gooseberries 5 by 4 feet, and Quinces and Dwarf Pear 10 or 12 feet.

It is a question of opinion whether trees should be planted in the Spring or Fall. Both seasons have their peculiar advantage. So far South there is little risk of injury from severe cold. Fall planted trees are sure to make better growth than those planted in the Spring, as new roots are emitted and the earth becomes firmly packed. Fall planting is done at a comparatively leisure season, and success is always largely dependent on perfect work. Purchases in the Fall are from full stocks, and more likely to be exactly as ordered.

NOTICE TO PURCHASERS.

Our terms are cash, unless otherwise agreed. At the price named, five will be furnished at half the cost of ten; fifty at half the cost of one hundred, etc. We pack trees in such a manner that they may reach the purchaser in good order, charging for time and material only. Please instruct how trees are to be marked and forwarded. We deliver all goods free to carriers in Lexington, taking receipts therefor, after which our responsibility ceases. Mail wagons and stage lines issue no receipts for goods entrusted to them, and we assume no responsibility in the event of loss when instructed to send over such lines. Small bales, boxes and parcels are best sent by express. We do not grow plants to be sent by mail, our soil not favoring the production of such light stock. The selection of varieties must be in part left to us; but if we are especially instructed to send only as *named*, the order will be filled in so far as we have the stock wanted. On lists without privilege please state if smaller or larger sizes than the ones named may be used. We wish to give perfect satisfaction in our dealings, and would rather return an order untouched than fill it unacceptably to the purchaser.

The propagation of all stock is under our immediate charge, and is true to name. We have always been and are ever ready to correct cheerfully all errors and shortcomings, and it is with pride that we say that in a sale of many millions of trees and vines, covering many years, not half a dozen complaints have been made of inaccuracy in labeling. Errors may and do occur, but they are not in the line of our interest.

We do not guarantee trees or plants purchased of us to live. No stock of doubtful vitality will be sent out, and we cannot insure against adverse seasons or the neglect of the purchaser in receiving promptly and planting with care, such as will insure success. Any loss that may occur through our fault will of course be adjusted at once and satisfactorily.

We cannot fill long lists of trees, etc., one of a kind, except at the price of a single tree. All letters answered at once when received, but write name and post office address plainly. Please order early, while stock is complete. We do not require trees to be taken when ordered; but will save them and send when wanted. The price list of this catalogue abrogates the prices of all previous issues. Money may be safely sent by check, post office order or registered letter. Our nursery is two miles north of Lexington, near the Georgetown pike, and on the Cincinnati Southern Railroad. Hillenmeyer Station, at which all accommodation trains stop, is but five minutes' walk from our grounds. We have no office in Lexington, and during the busy season can be found only at our packing grounds and plant rooms. A small Fruit Catalogue will be published in February and sent to patrons when issued, and others on application. We cordially invite the personal inspection of tree buyers and planters.

Address all communications to

H. F. HILLENMEYER,

Lexington, Ky.

Apples.

The Apple is the first fruit, both in commercial value and general adaption. By a judicious selection of varieties, a supply of this indispensable fruit may be had almost during the entire year. We are south of the latitude in which the apple attains its best estate, and few of the fine Northern varieties attain the same degree of perfection here. Northern winter apples ripen here during autumn. The apple requires four to eight years to attain the age of fruitage, and with ordinary care will reach an age of forty or more years. The second or third year after planting, the trees should be examined, and crowded or crossed branches should in March be removed. If this work is begun early, and a little pruning done each year, the trees will develop into handsome shape, and the necessity of removing large branches will never arise. The head of an apple tree should be kept open, and in the early state of development, when a pocket knife will suffice, is the time to determine its character. The saw and hatchet are the instruments of a heroic treatment under which many trees are permanently injured. Determine the character of head while it is small, and in the day of fruitage it will need little attention. It is unfortunate that many orchards are neglected until the trees are a mass of tangled branches. The pruning of such should be done from February to April 1st. Branches more than two inches in diameter are removed at great hazard. The pruning of badly neglected orchards should not be completed in one year, but by degrees. Large wounds should be covered with thick lead paint.

The diseases and injuries to which the apple tree is most liable are stem borers of various kinds, blight aphides, cancre and the gnawing of rabbits. We have found whitewash applied in May to the stems of young trees a preventative of the borer, and greatly beneficial otherwise. Twig blight and aphides seldom inflict serious harm, and cancre follows injury sustained by severe frost or the removal of large branches. No remedy can be applied when the injury arises from the first cause, and the correct treatment for the second has been indicated.

During periods of deep snow-fall rabbits frequently destroy young orchards by gnawing the bark near the snow line. If the stems are protected with straw, hemp or corn stalks, they will escape injury. To rub the stems with flesh, such as liver or other animal offal, is also an effectual remedy. The fruit of the apple tree is also liable to loss from many causes. The curculio and the codling moth are ever present enemies, and insidious fungous growths attack the young fruit when atmospheric conditions

favor these low forms of life. Intense heat and protracted drought are also frequent sources of failure. Indeed, premature ripening and falling off in consequence thereof is the greatest of all the causes of the failure of winter apples in our State. Our dry and warm autumns ripen varieties in September that should only be gathered in November.

It has been our unvarying custom to gather apples when ripe, whether it be the conventional season or not. Our crop has in the main, for many years, been gathered before September 15th, and there has rarely been difficulty in keeping it until May following.

Many excellent apples—Northern Spy, Rambo, Fallawater, Nonsuch, Bellflower, Milam, Grimes' Golden, etc., etc.—are a total loss, falling from the trees early in September. If these be carefully gathered and packed in sugar or other barrels, and the heads closely fitted, they will keep a long time. Early gathered apples must be free from all decay and bruises.

Spraying apples to destroy the curculio and codling moth has been thoroughly tried here with the most gratifying results. Our trees are twice sprayed early in May with water, in which is mixed one-quarter of a pound of London purple to fifty gallons. A tree may be sprayed in two or three minutes, and we have no hesitancy in advising our friends to procure an outfit. By applying to the Department of Agriculture, Washington, D. C., the latest researches in the treatment of insects and fungus pests may be had free of cost in condensed pamphlet form.

It is customary to seed the apple orchard with grass, but by occasional cultivation after the trees are in bearing and by timely care in pruning we are satisfied that half the climatic and insect cause of injury would be removed, and that the fruit would be larger and better, and the yield far more certain.

Apple trees at a given age vary materially in size, some kinds being of more vigorous growth than others. Our stock is unusually full this year and of fine quality.

SUMMER APPLES.

✓ **ASTRACHAN**—Red. A fine, showy apple; tart, rich and good; ripens over a long period. One of the best early apples.

✓ **CHENANGO STRAWBERRY**—Medium; striped with red. A handsome and good apple; tree hardy and productive. August.

✓ **EARLY HARVEST**—Medium size; yellow, flesh white, juicy and aromatic; bears regularly and is one of the most desirable of early apples.

✓ **EARLY PENNOCK**—A popular apple; large medium and of good quality.

✓ **EARLY TRANSPARENT**—(From Russia). Resembles Early Harvest; rather earlier, but not so good. This is of value.

✓ **FOURTH OF JULY**—Medium; whitish yellow, with red. An excellent early apple; productive and showy.

✓ **GRAVENSTEIN**—A famous old German apple, that is equally good here; large, red and yellow. August.

✓ **GOLDEN SWEET**—A very popular early sweet apple; excellent for baking; fine size. August.

✓ **GARDEN WALK**—A superb new apple raised by Mr. James Garrett, of Woodford County; size large medium, pale, greenish yellow, striped with red; superb in quality. August and September.

✓ **LEXINGTON**—Medium, with two shades of red; good quality; trees hardy and productive. July 20th.

MAIDEN'S BLUSH—Large, waxy yellow, with bright blush, flesh firm, fine-grained and aromatic; tree a regular bearer and most valuable in every way. August.

OLDENBURGH—Medium, yellow, deeply splashed with red. A regular bearer.

PRIMATE—Medium, yellow, with blush; fine quality. August.

WILLIAM'S FAVORITE—Large medium, red mostly, flesh white, juicy and highly flavored. August.

✓ **RED JUNE**—A very popular early apple, almost as well known as Early Harvest, and of same season; medium size.

✓ **SUMMER PEARMAN**—Best of all in quality; medium, red with russet; moderately productive. Tree small from nursery, but vigorous in orchard.

AUTUMN APPLES.

✓ **BAILEY SWEET**—A large, fine apple, red mostly, quality good; very productive. September.

✓ **FALL PIPPIN**—Large, rich yellow; one of the very best for drying; tree vigorous; a moderate but regular bearer.

✓ **FALLAWATER**—Large, pale yellow when fully ripe; flesh soft, rather coarse, tender and pleasant; tree vigorous, very productive, and one of the most valuable late autumn or early winter kinds.

✓ **GRIMES' GOLDEN**—Medium to large, deep yellow. This is a standard of excellence among apples. The tree is hardy, vigorous and productive. We prize this variety very highly. October to December.

✓ HUBBARDSTON'S NONSUCH—A superb October apple; yellow, deeply shaded with red; vigorous and productive.

✓ NORTHERN SPY—A large apple of the highest quality; well known. September and October.

✓ RAMBO—Medium size, yellow, striped with red; flesh white, crisp and aromatic. Best on warm soils.

✓ TWENTY-OUNCE—Very large, pale yellow. Tree healthy and productive.

✓ WINE APPLE—Large, pale red with yellow. A superb apple that succeeds well here.

WINTER APPLES.

✓ BALDWIN—A superb market apple that succeeds only moderately here. Bright red; large and good. December.

? BOSWORTH—A fine red winter apple, large, showy and most reliable; very late keeper. Named for Mr. Benj. Bosworth, Fort Springs, Ky.

BEN DAVIS—Large, handsome red; fruit very uniform in size; tree hardy, vigorous and productive; blooms late; a valuable apple for market.

BELLFLOWER—A large yellow apple of superior quality; well known. Not reliable.

✓ BLACK TWIG—TWITTY—KINNAIRD'S CHOICE—Seedlings of Wine Sap from Tennessee, and likely to prove of value. Very vigorous here; quality best.

✓ CLAYTON—A superb winter fruit, much resembling Red Pearmain. New and very promising.

✓ DR. WALKER—A seedling of the Janet; larger and brighter in fruit and of more vigorous growth. This variety promises to be of great value.

HUNTSMAN, MO., PIPPIN—Both popular West; fine size and quality, worthy of trial here. Trees healthy, strong growers.

JONATHAN—Medium size, red; fine quality; quite reliable. Early winter.

KING (Thompkins Co.)—A magnificent large red apple of the highest quality; tree a handsome grower, but not very hardy.

LADY APPLE—Small, waxy, transparent, with crimson cheek, superb quality and very late keeper. Tree hardy and productive.

✓ LIMBERTWIG—Resembles Janet; larger and brighter, but not equal in quality; tree vigorous, hardy and productive, a very late keeper.

✓ LAWVER—We hope that all is true that has been said of this. Tree vigorous; but not fruited here.

- MILAM—Medium size, red, of excellent quality; a regular and abundant bearer. December.
- NICKAJACK—A very popular Southern apple of good quality, and a late keeper.
- PARADISE SWEET—Yellow, with blush, medium size, good in quality, and a late keeper.
- ROMANITE (Small)—An old and very reliable late-keeping winter apple. Fruit medium size; good quality; deep red; tree an abundant and regular bearer.
- RAWLES' JANET—This has been the most popular apple West and South. Size medium; mostly red; flesh firm, crisp and sprightly; tree blooms late. Has rotted very badly in late years before maturity.
- RED PEARMAIN—An old and much esteemed variety; hardy, prolific and reliable.
- ROME BEAUTY—Large, with two shades of red; flesh firm, white and pleasant; tree most productive, not hardy, a late keeper.
- RUSSETT (Perry)—Large, yellow, very crisp and good. Unreliable.
- SMITH'S CIDER—Large, striped with yellow and red; flesh tender, juicy and crisp; valuable for market or home use; reliable as Ben Davis. January.
- STARK—Red and yellow; flesh firm, crisp and good, tree hardy and vigorous; growing in favor for its many excellencies. One of the best late keepers.
- TALMAN'S SWEET—Medium, pale yellow; one of the very best late sweet apples.
- VANDIVERE—Superb quality, fine size; tree vigorous and productive; falls badly. Early winter.
- WHITE PIPPIN—A fine, large fruit, the very best for canning; flesh white, juicy and aromatic. Quite reliable.
- WAGGENER—Medium to large; yellow, with bright blush; flesh juicy and fine grained.
- WHITE WINTER PEARMAIN—A very popular apple, large pale yellow with blush; a late keeper. Not so reliable as West.
- WINE SAP—Medium dark red, flesh yellow, tender, crisp and spicy; tree a small, tangled, irregular grower. Enormously productive and valuable on warm, rich soils.

✓ WINTER RAMBO—In size and appearance as Fall Rambo, but a late keeper; superb quality.

CRABS.

✓ HEWES' CRAB—Small, striped, enormously productive; very tart, making a dry cider unsurpassed for spice and piquancy; tree small and compact.

✓ HYSLOP—Large, dark red; tree vigorous and productive.

✓ MONTREAL BEAUTY—Most beautiful of crabs; waxy, yellow and carmine.

✓ RED CRAB—An improvement on Hewes' Crab in point of size and vigor of tree. Valuable for cider.

✓ SIBERIAN—Red and yellow. These apples are used only for preserving; size small; stem as long as that of a cherry; color waxy, with carmine blush.

✓ TRASCENDENT—An early ripening variety.

Peaches.

This admirable fruit attains its highest excellence on the dry soil of Central Kentucky. To be successful in the raising of the peach, it is necessary that the land be dry and the site elevated, as the fruit buds are more liable to loss on low ground from winter or late spring frosts. The trees should be annually pruned in November or March, *after they are in full fruitage*. This consists in the removal of three-fourths of the current growth. The borer found under the bark and just beneath the surface of the earth, must be faithfully removed. If this be done in October, the worms, still on the outside, may be easily removed by scraping the bark. Moist weather is most favorable for this work.

When but few peach trees are kept, perhaps the most perfect remedy for the borer is to make a slight cavity around the collar of the tree at the same season as advised above, and apply from a quart to a gallon of hot water, dependent on the size of the tree. The water may be nearly boiling, and the treatment is promotive of health.

✓ ALEXANDER—This is certainly one of the very best of extra earlies, ripening about July 1st. The fruit is of moderate size and excellent quality. Of beautiful color, it neither rots on the trees nor is too soft for market. Tree very hardy and productive, but does best on poor soil.

✓ CRAWFORD'S EARLY—This noble variety in thrift, productiveness, large size and general excellence, is unsurpassed among yellow-fleshed peaches; freestone. September.

✓ EARLY RIVERS—This is a superb early peach of large size and fine quality. Color straw, with blush; flesh very soft and melting. July 15.

✓ GEORGE FOURTH—Large freestone, ripening after Early York; unsurpassed in size, quality and appearance; desirable.

✓ HALE'S EARLY—Medium size, deep red; tree vigorous, very hardy and productive. July 20th.

HEATH CLING—This best known of peaches is unequaled among clingstones; size large; pale with occasional blush, flesh firm, crisp, very white, without a trace of red; stone small and flattened. Oct. 1st.

LARGE EARLY YORK—Medium, pale, mingled with red; superlative in quality. If confined to the planting of one tree, this would be our choice. August 1st to 10th.

✓ LAGRANGE—A very large, pale freestone, with bright blush; flesh very white to the stone. Best on dry, poor soil.

✓ LEMON CLING—Best of yellow clings; large, beautiful and good; productive.

LARGE WHITE CLING—This superb peach, cream color, with bright blush, is unrivalled among clingstones. September 1st.

✓ OLDMIXON CLING—Resembles the large White Cling closely.

✓ OLDMIXON FREE—Large red; flesh firm and rich; stone small; exceedingly valuable for home use or market. September 20th.

✓ PICQUET LATE—Fully as large as Crawford's Late, and superior in every way to that popular variety. September 20th.

✓ SMOCK FREE—Large, yellow and red; an old and popular variety. October. Not so good as the preceding.

✓ STUMP THE WORLD—A magnificent peach, very large, pale, with light red; flesh melting, rich and aromatic.

✓ In addition to the appended list, we have more or less of Henrietta, Chair's Choice, Elberta, Wonderful, etc., etc., and a few hundred seedling raised from mixed seeds of Alexander, Early York, George Fourth, Large White Cling and Crawford. The seeds were from the choicest fruit, and as we have application each year for seedlings for their supposed hardihood, a larger proportion of meritorious trees may be expected than is usual among seedlings.

Pears.

The ripening of this fruit, like the apple, extends over a long period. The earliest ripen the last of June, and winter pears will keep until February. Our bright, dry, warm autumns, however, are not favorable to the winter pear, for, like many apples, they ripen very much out of season under such circumstances. Standard pear trees are worked on seedlings, and are best suited for the orchard.

The stems of young trees should be wrapped with straw for several years. Pears are greatly improved by being gathered a few days before maturity, and then house-ripened in a dry, airy room.

The pear generally fruits in four or five years after planting, and is nearly as reliable as the apple. The tree is entirely hardy, and is healthy save in being liable to blight.

Dwarf pears are grown on quince roots, and are more especially adapted to the garden than standards. If planted on good ground, ten or twelve feet apart, and manured liberally, they will yield very fine fruit in three years, and prove a source of much pleasure.

Our stock of standards is larger than ever before and of extra quality.

BARTLETT—Best known of pears; very productive; large, yellow, rich and melting.

BEURRE D'ANJOU—Fine, large pear; flesh fine grained and vinous; tree hardy and productive. October.

BRANDYWINE—Medium to large; color russet and green; tree hardy and productive. August.

CLAPP'S FAVORITE—A seedling of the Bartlett, but larger, earlier and as valuable; tree vigorous and productive.

DUCHESS D'ANGOULEME—Very large, yellow and russet; best as a dwarf. October.

FLEMISH BEAUTY—A large, showy pear, most valuable for market. September.

GARBER—A hybrid ripening intermediate between Keiffer and Le Conte. Very promising indeed.

HOWELL—Large, yellow, of fine quality, ripening September 1st; an early and abundant bearer.

IDAHO—Well advertised, but certainly not blight proof. One year trees only.

- ✓ **KEIFFER**—A hybrid pear, ripening in October and November; tree vigorous, hardy and productive.
- ✓ **LECONTE**—Tree hardy, most vigorous and productive. September. Garber and Keiffer has thus far proved exempt from blight, and this class of pears, for their free growth and early and abundant bearing, with very fairly good quality, are each year growing in favor.
- ✓ **LAWSON**—An extra early pear of fine color and poor quality; valuable for market.
- ✓ **LAWRENCE**—Most valuable of winter pears; medium size; flesh melting and rich; tree hardy, productive and an early bearer.
- ✓ **SECKLE**—A well-known, small, russety fruit; the most exquisite of pears; of slow growth in nursery row. September.
- ✓ **SHELDON**—A noble American pear, worthy of a place in the smallest collection. September.
- ✓ **TYSON**—Medium, bright yellow, with red; flesh very sweet and aromatic.

Cherries.

This first fruit of the season succeeds best on dry, gravelly or stony soil. Heart or Bigarreau cherries are liable to injury from extremes of temperature, and if planted on our rich soils make such luxuriant growth that they will surely winter-kill. If planted in sheltered places, among trees and buildings, and not stimulated to luxuriance, if the heads are kept low and the stems wrapped with straw, these dangers are greatly lessened. Wet soils are surely fatal; even on dry ground, in nursery row after protracted rain, great loss frequently occurs during the season of rapid growth.

The fruit bud of the cherry is more hardy than that of the peach, and if this tree is consistently managed, a full or partial crop may be had almost every year. Cherries of the Early May type are entirely hardy, and while not equal to the others in use from hand, except if permitted to hang until thoroughly ripe, are very desirable, being exempt from most dangers that render the soft-wood kinds unsatisfactory. These are indicated by a star.

- ✓ **BLACK TARTARIAN**—Best known and most valuable of Heart cherries; fruit large, dark, half-tender, rich and pleasant; tree hardy, vigorous and productive.

✓ **BIGARREAU GRAFION**—Yellow Spanish; large, pale yellow, with red cheek; flesh firm, crisp and juicy; tree vigorous, hardy and productive.

✓ **BELLE MAGNIFIQUE**—Large, bright red, late; moderately vigorous and productive.

✓ ***CARNATION**—Large, light red, esteemed for preserving; tree vigorous; moderately productive.

✓ **DOWNER'S LATE**—Medium, light red; flesh soft, juicy and pleasant; late.

✓ ***EARLY MAY**—Known also as May Cluster and Early Richmond; is a tree of small growth; an early and most abundant bearer, and unexcelled for all culinary purposes. It is fully as hardy as the apple, and rarely fails to yield a full or partial crop.

✓ ***EMPRESS EUGENIE**—A small growing tree, very productive and hardy.

✓ **GOV. WOOD**—The best raised by Dr. Kirtland; large, light, with blush; tree vigorous and great bearer; very popular.

✓ ***MAY DUKE**—This popular variety has all the excellencies of Early May, and is of more vigorous habit.

✓ **NAPOLEON BIGARREAU**—Very large, waxy, with carmine blush; flesh firm, rich and delicious; tree symmetrical and productive.

✓ **OHIO BEAUTY**—Large, pale and dark red; tender, juicy and good, quite early.

✓ **ROCKPORT**—Large, pale and red mingled; tree upright and symmetrical.

✓ ***REINE HORTENSE**—A late, large kind, and best of its class for use from hand.

Plums.

The plum succeeds best on heavy clay loams, making a compact tree, rather smaller than the peach. The plum is quite hardy, and will endure very nearly as much frost as the apple without injury. Trees planted where the ground is hard trodden yield the best, but they should be cultivated until well established. The American varieties, Wild Goose, Miner, etc., though inferior to the European type, are more hardy, healthy and productive, and generally escape the curculio with but little injury. The plum, when planted in city yards, is notably exempt from the attack of the curculio, the greatest enemy of the plum. We have observed plum trees in town inclosures bearing magnificently year after year, and recommend with confidence this tree for such locations. The tree itself is com-

pect, with clean, healthy foliage, and when covered with fruit is a striking object on the lawn or in the garden.

The European plum is of superior quality and value, but is prone to the black knot. If the trees are closely watched in June and July, and the excrescences cut away while soft, the remedy is perfect. To stimulate the trees to a vigorous annual growth is also excellent remedial practice, the disease seldom attacking trees in full thrift. The curculio frequently destroys every plum, and then almost entirely disappears. In the past, the insertion in the stem of an iron pin, upon which to strike with a hammer, has shown good results. The injury inflicted by the curculio is done from early May to mid June, and the jarring of the trees was labor well repaid. The curculio may be effectually suppressed by spraying, but the solution should be a weak one; plum and peach foliage being sensitive to arsenical poisons.

American plums yield best when several varieties are planted near each other, the flowers being thus more surely fertilized.

✓ BRADSHAW—A popular, large and good plum; tree vigorous, hardy and productive.

✓ COE'S GOLDEN—Late, large and good; tree moderately vigorous and productive.

✓ DAMSON—A variety too well known to need description; succeeds well almost everywhere.

✓ DOUANE'S PURPLE—Very large; purple with white plume; flesh soft, sprightly and adheres to the stone. August 1.

✓ GREEN GAGE (Reine Claude)—Medium size; yellow, with plume; flesh soft, rich and aromatic; one of the best in every way; tree moderate grower. August.

✓ IMPERIAL GAGE—As above, but nearly twice as large; not so rich; color bright yellow. September.

✓ LOMBARD—Medium size; violet red; flesh yellow, brisk and pleasant. Best.

✓ MINER—An improved Chickasaw; quite late and good quality; resists the curculio well.

✓ MARIANNA—Not so early as Wild Goose, but proves very desirable.

✓ POND'S SEEDLING—Large, red and yellow; tree vigorous and productive.

✓ WEAVER—One of the Iron Clads; fruit medium, purple and of good quality. September 15.

✓ WILD GOOSE—An improved Chickasaw; size medium, color deep red, with bright plume; quality medium to good. A most abundant bearer, of great value. July.

Quince.

The quince delights in a deep, retentive soil, and should be cultivated or mulched if fine fruit is expected.

The trees bear in three or four years, and are enduring, hardy and productive.

APPLE OR ORANGE—This is the most highly esteemed variety; productive of smooth and perfect fruit. September.

CHAMPION—More vigorous than above and later; not very productive here.

Apricots.

These ripen rather earlier than the peach, and partake in quality of both the peach and plum. The standard old kinds are of no value here, but the hope is entertained that the Russian type may prove more hardy and enduring. They are not yet tested here, and are sent out with many misgivings as to their value. Assorted varieties, 3-4 feet, 25c each.

Currants.

These succeed best on deep, cool soil, and should be planted four feet apart and kept well manured or mulched. A north wall is a favorable place for them. So far south they are prone to lose their foliage at mid-summer, and many desirable kinds north are valueless.

The fruit is highly esteemed for many culinary purposes, and will hang on the bushes for a month or more. The white varieties are of no value here. The old wood should be thinned occasionally. The summer just passed has not been favorable to these, being too warm.

BLACK NAPLES—The best black-fruited variety; very vigorous and quite productive; bunch small; flavor peculiar.

CHERRY—Very large; plant vigorous; not reliable.

FAY—Long cluster and of good quality; productive; moderately vigorous.

LA VERSAILLES—A foreign variety, of great beauty, size and productiveness.

RED DUTCH—The best of currants; very vigorous, hardy and enormously productive; the only variety on our grounds that retains its foliage

through periods of extreme heat or drought, and better for all purposes than any variety we have tested.

VICTORIA—A standard kind that succeeds fairly well here.

Gooseberries.

This fruit requires the same soil and management as above. The oldest wood should be annually removed to insure abundant crops of fine fruit. The foreign varieties are worthless in this warm climate. We have quite a variety of the newer kinds, but are not satisfied as yet of any marked advance.

AMERICAN—Vigorous; fruit medium size; enormously productive; one of the best.

HOUGHTON—Vigorous; very productive.

SHAKER—A very vigorous, hardy kind, and quite productive; late, with large fruit.

Strawberries.

Experience and observation have taught us that these do not succeed well planted in autumn. In earlier years we furnished plants at this season, but the loss to the purchaser was so certain that we do not care to furnish plants, save when the buyer is prepared to lose both his money and labor.

We have this year larger plantings than ever, but the prospect for a full stand of plants is not encouraging. A special catalogue will be issued in February.

Raspberries.

These succeed the strawberry in time of ripening, and are of easy growth, hardy, productive and indispensable. The plants should be set three feet apart in rows five feet asunder. In planting red varieties, draw a deep furrow, drop the plants, draw enough earth around to hold them in position, and then cast earth from both sides with plow, shortening the canes at top of ridge thus formed. The young canes will push from beneath the surface, and are frequently quite late in appearing. Black va-

rieties should also be planted in a deep furrow, to prevent the stools from being blown down in later years when heavy with fruit and foliage. The plants should, however, be covered only an inch or two at the time of setting, as the young canes do not penetrate readily any great depth of covering. The drills are leveled by subsequent cultivation. When plants are set in the fall they should be amply covered, to prevent drawing by winter frost. Part of this covering should be removed at the approach of spring. Raspberries do best on moderately rich, moist soil. The raspberry is one of the most certain fruits that can be planted in this latitude. For many years we have had large stocks of bearing plants, and have uniformly gathered crops that paid handsomely. This fruit is not found in the majority of gardens, from a mistaken idea as to its proper management. It will not flourish in fence corners, unpruned, uncultivated and neglected. The planting should be done as indicated, and the succeeding year cultivation should be frequent, though quite shallow. After the first season's growth the young canes should in November or March be severely pruned, leaving the plants from twelve to eighteen inches high, shortening also the side branches to a length of six or eight inches. The second year there should be a very satisfactory crop of fruit and a great development in the size of the plant. The cultivation should be as the year preceding. Red varieties will sprout abundantly from every root; but unless all these are destroyed, save such canes as spring up within the stool itself, failure will ensue, as the plant is unable to support a crop of sprouts and perfect a crop of fruit. As the young canes reach a height of two feet they should be pinched in, thus causing short, stocky wood, well branched. Of late rust has attacked the black varieties in many localities, making them very unsatisfactory.

Immediately after the last gathering, the old canes are removed and the young ones thinned, five to ten being left dependent on the strength of the stool. It is indispensable to success that the cultivation be perfect: that the sprouts on red varieties be destroyed by hoe and plow; that the plants be pinched in to obviate the necessity of staking, and that the winter pruning be thorough, no lateral branch being more than a foot long.

✓ CAROLINE—Yellow raspberries are not desirable. This is one of the best, and we have plants of several other kinds, but do not recommend them except for amateur growers.

✓ GREGG—A late and large berry, black, firm, moderate in quality; ripening very uniformly. It is a good variety for market.

✓ MAMMOTH CLUSTER—An old reliable and very popular kind.

OHIO—An early black cap that in vigor and productiveness is almost unequaled. Fruit medium in size and of good quality.

QUEEN OF THE MARKET, OR CUTHERBERT—We find this fairly hardy, vigorous and productive. The fruit is large, late, rather dark, firm, but only moderate in quality. Its size, firmness and late ripening are valuable qualities in a market variety.

✓ SHAFER—The largest berry we have, and plant very vigorous and productive. Though of bad color; it has become very popular; late quality good; best for canning and jam.

✓ TURNER—Best of the red varieties; hardy; vigorous and most productive. Neither heat nor cold seems to injure the plant. Turner has no thorns. If the numerous sprouts that spring from the roots of bearing plants be hoed off as soon as they appear, nothing further in thrift, hardihood, quality or productiveness need be desired.

Asparagus,

Earliest of vegetables and should be in every garden. This vegetable is not as generally found in gardens as its merit deserves, from the prevalence of many erroneous impressions and practices in its management and culture.

The soil *must* be rich and should be dry and warm. A pit filled with manure as a bed for the plant is not needed. After a thorough preparation of the soil, draw three drills, eighteen inches apart, four or five inches wide and four inches deep. Spread the roots spider shaped in the drills, twelve or fourteen inches apart, and cover. No covering of manure is needed at this time. When the plants appear, cultivate and continue throughout the season to keep the earth clean and mellow. When the tops are frosted, mow and apply two or three inches of manure. Fork the bed lightly the following spring, continuing the culture as the year previous, and a similar application of manure in November each year thereafter. If the growth the first two seasons has been vigorous, the plants may be cut for the table slightly the third season.

If growing for market, the plants may be set in checks three feet apart each way. With more room for development, less manure will be required to produce fine stalks. The success of Asparagus culture is, however, dependent on the liberal application of manure annually, and the suppression of weeds during the period of growth.

The causes of failure are many. If the plants are set six or eight inches deep when received they will not penetrate the depth of covering, and will perish. A covering of manure on new-set plants is also generally fatal. To cut plants for the table the first year after planting is ruinous. To mow the tops of the plants while in growth is a prevailing custom, and it is the practice employed to destroy such plants as Ironweed, Poke, etc. It is needless to say such practice would kill outright any save a plant of vigorous constitution. To mow the tops when the seed ripens is also erroneously recommended. The seeds will not injure, and not one in a million will make a plant

A curbing of stone is a useless appendage to an Asparagus bed. Moles

do not injure *Asparagus* plants, and the curbing will not prevent their entrance.

Salt, liberally applied, renders the ground cold and clammy, and while its use is quite general, our experience does not indicate that the practice is a wise one.

The fertility of an *Asparagus* bed is greatly promotive of rank weeds. When the plants stand uniformly they will generally suppress such growth, but in vacancies they will spring up with great vigor, much to the injury of the bed, and must be removed. In late years, immediately after discontinuing the cutting of our beds for the table, we loosen the soil and cover with three or four inches of straw or hemp shives which effectually suppresses all growth of weeds, and is conservative of both moisture and fertility.

From many years of experience with various varieties we question whether there is a difference. Individual plants are superior to others, but the superiority cannot be perpetuated by seed, which is the universal method of propagation. This view is held by the best of botanists. Excellence is the result of liberal manuring and the selection of the heaviest and best seed. Our method of seed saving is peculiar, and in size and weight cannot be surpassed. The stock of plants on hand this year is very fine and large.

Rhubarb.

This, like the foregoing, is an indispensable early vegetable. The success of this plant is also dependent largely upon clean culture and abundant manure. The plants should be set three feet apart, with crown of plant three inches below the surface. The plants should be manured annually and heavily mulched with straw. A division of crowns every third year greatly increases the size of the stalks. When the stools are old the leaf-stems are numerous, but small; hence the necessity of division. Victoria is one of the best kinds, being unsurpassed in size of leaf-stem.

Grapes.

There is no branch of fruit-growing at present attracting so much attention in Kentucky as grape culture, and the reason is quite apparent. The grape-vine is very hardy, of easy growth, bears the third year after planting, and seldom fails to yield a full or partial crop. Aside from the certainty of the crop, the yielding per acre exceeds that of any other fruit, the apple not excepted, nor even the potato. Grapes, while hardy and prolific, are imperious in their demand for specific management. They should be planted seven or eight feet apart, each way, the plants being inclined at an angle of 45 degrees, and the roots covered six or eight inches. No stakes are needed the first year. The cultivation the first year and ever after should be frequent, but not deep. After the first season's growth the vines should be pruned in November, removing the entire growth save two or three buds. Stakes or a trellis made with wire and posts is needed to support the growth of the second season, each vine being limited to the making of but one, or if very vigorous, two canes. It is

imperative that all others be removed after danger of frost is past. At the end of the second year these canes must be shortened to four feet. The third year the vines will fruit, but they will require an annual pruning proportionate to the vigor of the plant. We rarely leave more than one-fourth of the wood of *current growth*, and if the plant lacks vigor the pruning should be more severe. Trimming and cultivation are the secrets of success in grape culture, and if neglected failure is certain.

It is unfortunate that when a vine is trellised to an old building it will yield abundantly and even very fine clusters without pruning. Such locations always engender the deposition of large quantities of nitre, which is extremely promotive of vigor in the grape vine. When pruning is named as a necessity to success, these vines are always cited as evidence of the incorrectness of the advice given. The peculiar circumstances of the location are the cause of the favorable result. It is the experience of vine dressers of all time and countries that the vine, save the Scuppernong, must be annually dressed.

Grape-vines of native stock are not subject to disease. The fruit when quite small is slightly liable to mildew in humid weather, and when half-grown under the same atmospheric conditions rot sometimes prevails, greatly lessening the yield. Both rot and mildew may be entirely prevented if the clusters are covered with paper sacks when the fruit is quite small.

Rot may be effectually controlled with any of the standard applications, and our vineyards, of many thousand vines were this year very satisfactorily protected at small cost. We used Eau Celeste and Bordeaux mixture.

Sacking grapes not only obviates disease, but greatly favors handsome development and high quality. Sacked grapes are not injured by bees, wasps, birds, etc., and will keep perfect on the vine until severe frosts.

We have protected grapes in this way, at very small cost, for several years, and recommend the practice with confidence. Grapes may be kept for several months if placed in thin layers in a dry, frost-free room.

We have been so greatly impressed with the value of the grape for general planting that great pains have been taken in collecting and testing new varieties and old ones not generally disseminated. After many years of patient care and observation we find that few varieties possess the qualities requisite for general planting, and to-day we have hundreds of dollars worth of high priced grape vines consigned to the brush heap, and as many more in our vineyard that will likely prove of as little value.

The demand from nine-tenth of our patrons is for vines that are hardy, healthy, vigorous and productive, and of quality that is fairly good. After a full test of hybrid vines—crosses between the vine of Europe and native types—we find them worthless. Many other varieties of purely native origin are not valuable. The list appended is short, but on our grounds they have stood the severest test of adverse condition incident to the climate of our State. Most of the varieties named are so vigorous in growth that the foliage in July may seem superabundant and injurious, inducing many cultivators to summer prune. This we have invariably found injurious to both fruit and the vigor of the vine. The better way is to pinch the ends of the cane as soon as it is indicated that the trellis is

insufficient to accommodate the unchecked growth. Pinching stops terminal growth and causes a sturdier habit of wood, superior for next year's fruiting, to the more attenuated wood of a vine whose growth has been at will. In addition to the list named, we can furnish wood or vines of perhaps one hundred other kinds.

- ✓ BRIGHTON—A superb red grape; hardy and moderately vigorous. In quality many have pronounced it superior to Delaware.
- ✓ CATAWBA—This old grape is unsurpassed in quality. It did poorly for a long time, but of late years our vines bore heavy crops of handsome fruit.
- ✓ CONCORD—This is undoubtedly the best grape we have for general planting. The vine is perfect in thrift, hardihood and productiveness. The bunch and berry are large and the flesh melting. The flavor is somewhat foxy, but when well ripened, especially in sacks, this grape, under our warm skies, is truly excellent.
- ✓ COTTAGE—A seedling of the above, closely resembling its parent, but ten days earlier. Not quite so good in bunch and quality, but one of the best early grapes in our collection and the hardiest.
- ✓ DELAWARE—This noble grape, in quality, is the best, and is perfect in color and beauty of cluster. The vine is hardy, but a very tardy grower. It must have rich soil and good culture.
- ✓ ELVIRA—Hardy and very productive. Bunch perfect; berry medium; flavor pure; ripening after Concord. This variety must be pruned very closely, or it will overbear, the fruit then being very insipid. White.
- ✓ IVES' SEEDLING—This vine is nearly as hardy as Concord; more vigorous and twice as productive. It is not as good a table grape, but the best wine grape for general planting yet introduced. Color black; bunch large and perfect; flavor rather foxy, but when fully ripe sweet and aromatic. It will hang on the vine until shriveled.
- ✓ MARTHA—This standard white grape is larger in bunch and berry than Elvira, and is an old favorite.
- ✓ NORTON'S VIRGINIA—This old and reliable grape is a general favorite. Rampant grower, ordinarily hardy and very productive. The bunch is perfect and large and berry small. It ripens late and will hang until frost falls. Should be pruned to canes rather than spurs.
- ✓ NOAH—This white grape will prove an acquisition. It is perfectly hardy, very healthy, and our vines have proved astonishingly productive. It ripens late. Bunch and berry resembles Martha. Quality moderate. Drops from cluster when ripe.
- ✓ NIAGARA—A fine white grape; plant vigorous, healthy and hardy. Cluster very fine. Should be pruned to canes rather than spurs.
- ✓ PERKINS—Vine hardy, vigorous and productive; cluster medium; berry large, dull red, very musky in flavor.
- ✓ POCKLINGTON—A seedling of the Concord, with large bunch and berry. This and Niagara are more prone to rot than many other standard kinds.
- ✓ WORDEN—Closely resembles Concord in leaf, wood and habit of vine. Ripens earlier, cluster and berry large; pulp soft and melting; vine hardy, vigorous and productive. This grape has come to stay.

✓ WYOMING—This new red variety is of good quality, with handsome clusters. Hardy, vigorous and productive; an acquisition.

✓ WOODRUFF—Another red variety, of which we hope great things. Quality moderate, but perfect in cluster and of beautiful color.

Persimmon.

This admirable native fruit has been very much neglected. The persimmon fruits quite early, and blooming very late, entirely escapes untimely frosts. The tree, yearly laden with its golden fruit, is highly ornamental on the lawn, and months after the orchard is brown and bare retains its fruit quite uninjured by midwinter frost.

The persimmon does not transplant kindly, even when nursery grown and should be set with extra care, protecting the stem with straw. Clay soils are most congenial to this tree.

Ornamental Trees.

Nothing adds so much to a home in monetary value and true comfort as a well planted lawn. A home without trees and shrubs is forlorn indeed. Tree planting on the lawn is not so generally satisfactory as in the orchard, from a neglect to have the holes large and well prepared, and from a failure to mulch or at least to suppress the grass immediately around the roots. To sod around a newly planted tree means failure. If the season is dry, complete loss is almost certain, and at best but feeble growth may be expected. For rapid growth and vigorous trees thorough planting and a perfect mulch are necessary. The branches of most lawn trees should be shortened one-half when set and the stems wrapped.

Our stock of both deciduous and evergreen ornamental trees is very full this year, and large planters using car lots will be favored with very low rates.

ASH (European)—Resembles the White Ash, but more rapid in growth and erect in habit. A fine stock this year.

CAROLINA POPLAR—A most rapid growing tree, resembling Lombardy Poplar, but with larger leaves and more open habit. Valuable for street planting and rapid effect. Very popular.

CATALPA SPECIOSA—A fine lawn tree, with large, dark foliage, more erect than the common Catalpa, blooming later, and very largely planted at present for timber.

CATALPA JAPAN—Resembles the above, but more erect; foliage darker, with very abundant flowers, very rapid while young.

CHESTNUT (American)—A well-known nut tree, of rapid growth and fine form. It establishes slowly.

ELM (Slippery)—Well known for the peculiarities of the inner bark. Tree of rapid growth, with large foliage.

ELM (Camperdown and Scotch)—Not so rapid in growth as the American, but more graceful and very pretty. Fine stock.

ENGLISH ALDER—A very pretty and rapid-growing tree, desirable on the lawn. Superior stock this year.

ENGLISH WALNUT—Well known as furnishing the imported walnut of

commerce. Tree hardy here and of rapid growth. Fruits moderately: Fine stock this year.

ENGLISH OAK—More rapid than native kinds.

HYDRANGEA—Hardy; a most desirable late-blooming shrub, bearing immense spikes of white flowers and remaining a conspicuous object on the lawn at a season when few shrubs are in bloom.

LABURNUM—A pretty, small-growing tree, with handsome flowers.

LARCH (European) A very graceful tree, with foliage resembling that of an evergreen. Fine Stock.

LIQUIDAMBER—Desirable for its gorgeous autumn foilage. Rapid growth.

LINDEN (European)—Broad-leaf and Red-twigged. These are graceful trees, rapid in growth, with smaller foliage and more erect in habit than the American.

LINDEN (American)—A noble native tree, of extremely rapid growth, with excellent habit and foliage. It is largely planted by apiarians, yielding an excellent article of honey.

MAPLE-SYCAMORE—An erect and vigorous tree, developing into perfect form; foliage dark. Desirable.

MAPLE (Norway)—This superb tree has all the beauty of the Sugar Maple, and is of much more rapid growth. With very dark foliage, expanding early and retained late, it forms a perfect globose head of unsurpassed beauty. We have a very large stock.

MAPLE (Silver)—The Silver Maple is unequalled for general lawn planting. Of easy and rapid growth, it requires but a few years to attain considerable size; is hardy, healthy and free of insects. It is first in expanding its leaves and retains them green as ivy until severe frost. Of these we have many thousands, and can furnish orders with choice trees, admirably rooted, being one or more times transplanted.

MAPLE (Sugar)—Of rather slow growth until established, but makes up for lost time afterward, developing into a majestic and symmetrical tree; earliest green in spring, and gorgeous with crimson and gold for a month or more in autumn. Our trees are nursery grown and extra good.

MAIDEN HAIR -Salisburyia—A superb Japan tree that is not well known, but will become popular. Stiff as an Oak, yet pendulous and graceful as a Willow or Birch.

MAGNOLIA—Accuminati—Well known for its stately habit, and beautiful flowers.

MOUNTAIN ASH—A tree of rapid growth, bearing large clusters of red berries until severe frost. Large stock.

*POPLAR-LOMBARDY—A very erect tree, of extraordinarily rapid growth, valuable on the lawn to produce striking contrasts.

PURPLE FRINGE OR SMOKE TREE—A popular shrub, much admired for its large clusters of flowers, which are very persistent, making it a striking object on the lawn.

RED BUD—A well-known tree, blooming very early in the spring, before the foliage expands.

WHITE FRINGE—A very ornamental shrub, bearing white racemes of flowers in June.

WHITE BIRCH (European)—A graceful tree, of neat habit; bark white; foliage dark and glossy.

WILLOW (Weeping)—A tree well known for its pendulous habit of growth.

YELLOW POPLAR (Tulip Tree)—A native tree of fine foliage and erect habit. It is very rapid in growth and remarkably free from disease. It bears tulip-shaped flowers; fine stock.

Hardy Flowering Shrubs.

We offer a fine stock of shrubs for the lawn, entirely hardy and of fine blooming size. The list consists mostly of Calycanthus, Deutzias, Burning Bush, Lilacs, Honeysuckles, Syringas, Weigelas, Tamerax, Snowballs, Spirias, Pyrus Japonica, etc., etc.

Evergreens.

Type of the immortal, and amid the desolation of winter, keeping bright the promise of life! These should be found on every lawn. They are easily transplanted as other trees, and are more rapid in growth than many. Most evergreens have soft fibrous roots, and if placed in position and earth shoveled over them without using the hand to lift and spread them into a plane of earth as the one occupied before digging, they are liable to fail. This is especially true of Hemlock, Arborvitæ and Juniper. A superb stock this year, and special rates on all large orders and car lots.

NORWAY SPRUCE—The most popular evergreen west; hardy, healthy, of rapid growth, and fine form and foliage.

HEMLOCK—This native tree, when well grown, is unsurpassed among evergreens. It is erect, yet pendant, of symmetrical contour, and delicate foliage. Succeeds here admirably.

PINE (Austrian)—A tree of stout and vigorous habit; very broad in growth; foliage of the darkest hue.

SILVER FIR—An erect tree, of rapid growth; foliage of bright green above and silvery beneath.

JUNIPER (Irish)—The best of our erect evergreen shrubs, attaining a height of ten feet. In favorable soil it makes a magnificent pyramid of brightest green.

ARBORVITÆ (American)—A pretty tree of rather dwarf growth. It is much used for screens and ornamental hedges. Our stock this year is unusually fine.

ARBORVITÆ (Chinese and Siberian)—Darker in foliage and more compact than American.

WHITE PINE—This well known tree is most rapid in growth, developing into a majestic and symmetrical form, with foliage perfect in color.

RETINOSPORA (in variety)—Dwarf shrubs, with dense, slender, feathery branches; very ornamental; fine stock. These are not as popular as their merit deserves.



